

# On the parity of ranks of Selmer groups II

Jan Nekovář

11 November 2000

Let

$$E : y^2 = x^3 + Ax + B \quad (A, B \in \mathbf{Q})$$

be an elliptic curve over  $\mathbf{Q}$  of conductor  $N$ . Thanks to the work of Wiles and his followers [BCDT] we know that  $E$  is modular, i.e. there exists a non-constant map  $\pi : X_0(N) \rightarrow E$  defined over  $\mathbf{Q}$  and

$$L(E, s) = \sum_{n=1}^{\infty} a_n n^{-s} = L(f, s)$$

for a normalized newform  $f \in S_2(\Gamma_0(N))$ .

For a large class of number fields  $F$  (which includes all solvable Galois extensions of  $\mathbf{Q}$ ), this is known to imply that the  $L$ -function  $L(E/F, s)$  has a holomorphic continuation to  $\mathbf{C}$  and a functional equation relating the values at  $s$  and  $2 - s$ . For such  $F$ , denote by

$$r_{an}(E/F) := \text{ord}_{s=1} L(E/F, s)$$

the *analytic rank* of  $E$  over  $F$ .

Over an arbitrary number field  $F$ , the  $m$ -descent on  $E$  gives rise (for every integer  $m \geq 1$ ) to the Selmer group  $S(E/F, m)$  sitting in the standard exact sequence

$$0 \rightarrow E(F) \otimes \mathbf{Z}/m\mathbf{Z} \rightarrow S(E/F, m) \rightarrow \text{III}(E/F)[m] \rightarrow 0.$$

Fix a prime number  $p$  and put

$$\begin{aligned} S(E/F) &= S_p(E/F) = \varinjlim_n S(E/F, p^n) \\ X(E/F) &= X_p(E/F) = \varprojlim_n S(E/F, p^n) \\ s_p(E/F) &= \text{corank}_{\mathbf{Z}_p} S_p(E/F) = \text{rk}_{\mathbf{Z}_p} X_p(E/F). \end{aligned}$$

The conjecture of Birch and Swinnerton-Dyer predicts that

$$(BSD) \quad r_{an}(E/\mathbf{Q}) \stackrel{?}{=} \text{rk}_{\mathbf{Z}} E(\mathbf{Q}).$$

As in [NePl], we are interested in a rather weak consequence of (BSD) (and the conjectural finiteness of the Tate-Šafarevič group), namely the

$$\textbf{Parity conjecture for Selmer groups:} \quad r_{an}(E/\mathbf{Q}) \stackrel{?}{\equiv} s_p(E/\mathbf{Q}) \pmod{2}.$$

Our main result is the following

**Theorem A.** *Let  $E$  be an elliptic curve over  $\mathbf{Q}$  with good ordinary reduction at  $p$ . Then the parity conjecture*

$$r_{an}(E/\mathbf{Q}) \equiv s_p(E/\mathbf{Q}) \pmod{2}$$

*holds.*

See [NePl] for a discussion of earlier results in this direction. Our method of proof is similar to that in [NePl]; the only difference is that we use anticyclotomic deformations instead of Hida families. A recently proved conjecture of Mazur ([Maz2], [Co], [Va2]) on non-vanishing of Heegner points in anticyclotomic  $\mathbf{Z}_p$ -extensions plays the role of Greenberg's conjecture assumed in [NePl].

## 1. Heegner points

In this section we recall the basic setup of Heegner points on  $E$  (see [Gro] for a more detailed account).

**1.1** Let  $K = \mathbf{Q}(\sqrt{D})$  be an imaginary quadratic field of discriminant  $D < 0$ . We assume that  $K$  satisfies the following “Heegner condition” of Birch [Bi]:

(Heeg) Every prime  $q|N$  splits in  $K$ .

Under this assumption there exists an ideal  $\mathcal{N} \subset \mathcal{O}_K$  such that  $\mathcal{O}_K/\mathcal{N} \xrightarrow{\sim} \mathbf{Z}/N\mathbf{Z}$  (of course,  $\mathcal{N}$  is not unique; we choose one).

**1.2** For every integer  $c \geq 1$  denote by  $\mathcal{O}_c = \mathbf{Z} + c\mathcal{O}_K$  the unique order of  $\mathcal{O}_K$  of conductor  $c$ . If  $(c, N) = 1$ , then

$$\mathcal{N}_c := \mathcal{O}_c \cap \mathcal{N}$$

is an invertible ideal in  $\mathcal{O}_c$  satisfying  $\mathcal{O}_c/\mathcal{N}_c \xrightarrow{\sim} \mathbf{Z}/N\mathbf{Z}$  (and hence also  $\mathcal{N}_c^{-1}/\mathcal{O}_c \xrightarrow{\sim} \mathbf{Z}/N\mathbf{Z}$ , since  $\mathcal{N}_c$  is invertible). The cyclic  $N$ -isogeny

$$[\mathbf{C}/\mathcal{O}_c \longrightarrow \mathbf{C}/\mathcal{N}_c^{-1}]$$

induced by the identity map on  $\mathbf{C}$  defines a non-cuspidal point on the modular curve  $X_0(N)$ , which is defined over  $H_c$ , the ring class field of conductor  $c$  over  $K$ . The image of this point under the fixed modular parametrization  $\pi : X_0(N) \longrightarrow E$  will be denoted by  $\bar{x}_c \in E(H_c)$  – this is a *Heegner point of conductor  $c$  on  $E$* .

**1.3** From now on, we consider only conductors of the form  $c = p^n$  for a fixed prime number  $p$ . The field  $H_{p^\infty} = \bigcup H_{p^n}$  then contains the *anticyclotomic  $\mathbf{Z}_p$ -extension*  $K_\infty$  of  $K$ , which can be characterized by the following properties:  $K_\infty = \bigcup K_n$ ,  $G(K_n/K) \xrightarrow{\sim} \mathbf{Z}/p^n\mathbf{Z}$ ,  $K_n/\mathbf{Q}$  is a Galois extension with  $G(K_n/\mathbf{Q}) \xrightarrow{\sim} D_{2p^n}$  (the dihedral group). There is  $n_0 \geq 0$  such that

$$H_{p^{n+1}} \cap K_\infty = K_{n+n_0} \quad (n \geq 0).$$

We use the following standard notation:

$$\Gamma = G(K_\infty/K), \quad \Gamma_n = G(K_\infty/K_n), \quad \Lambda = \mathbf{Z}_p[[\Gamma]].$$

Fix an isomorphism  $\Gamma \xrightarrow{\sim} \mathbf{Z}_p$  (i.e. fix a topological generator  $\gamma$  of  $\Gamma$ ).

From now on, we assume that the following condition of “good reduction” is satisfied:

(G)  $p \nmid N$ .

This assumption implies that the Heegner points

$$\bar{x}_{p^{n+1}} \in E(H_{p^{n+1}})$$

are defined. We put

$$x_{n+n_0} := \mathrm{Tr}_{H_{p^{n+1}}/K_{n+n_0}}(\bar{x}_{p^{n+1}}) = \mathrm{Tr}_{H_{p^\infty}/K_\infty}(\bar{x}_{p^{n+1}}) \in E(K_{n+n_0}) \quad (n \geq 0).$$

**1.4** In his lecture at ICM 1983, Mazur formulated (among others) the following

**Conjecture [Ma2].**  $(\exists n \geq n_0) \quad x_n \notin E(K_n)_{\text{tors}}.$

This conjecture was recently proved by Vatsal [Va2] under the assumptions that  $|D|$  is prime and  $p$  does not divide the class number of  $K$ , and by Cornut [Co] under the assumption (G). Both [Co] and [Va2] build upon an earlier work [Va1] of Vatsal.

**1.5** The fundamental distribution relation for Heegner points ([PR], Lemma 2, p.432) states that

$$\text{Tr}_{K_{n+1}/K_n}(x_{n+1}) = a_p x_n - x_{n-1} \quad (n \geq n_0 + 1).$$

The assumption (G) is equivalent to  $E$  having good reduction at  $p$ . From now on, we assume in addition that  $E$  has *ordinary* reduction at  $p$ , which is equivalent to

$$(Ord) \quad p \nmid a_p.$$

This assumption implies that the local Euler factor of  $E$  at  $p$  factorizes as

$$1 - a_p X + pX^2 = (1 - \alpha X)(1 - \beta X),$$

where  $\alpha, \beta \in \mathbf{Z}_p$  satisfy  $\text{ord}_p(\alpha) = 0$ ,  $\text{ord}_p(\beta) = 1$ .

Define, for  $n \geq n_0 + 1$ ,

$$y_n = x_n \otimes \alpha^{1-n} - x_{n-1} \otimes \alpha^{-n} \in E(K_n) \otimes \mathbf{Z}_p.$$

Then

$$\text{Tr}_{K_{n+1}/K_n}(y_{n+1}) = y_n \quad (n \geq n_0 + 1),$$

i.e.  $y = (y_n)$  is an element of the projective limit

$$\varprojlim_{n \geq n_0} (E(K_n) \otimes \mathbf{Z}_p) = \varprojlim_n (E(K_n) \otimes \mathbf{Z}_p) \subseteq \varprojlim_n X(E/K_n) =: X_\infty.$$

We shall also be interested in the inductive limit

$$S_\infty = \varinjlim_n S(E/K_n).$$

Both  $X_\infty$  and  $S_\infty$  are  $\Lambda$ -modules (of finite and co-finite type, respectively).

## 2. Iwasawa theory

In this section we recall basic results of Iwasawa theory of elliptic curves relating the  $\Lambda$ -modules  $X_\infty$  and  $S_\infty$  to Selmer groups over the fields  $K_n$ . The assumptions (Heeg), (G) and (Ord) are in force.

**Lemma 2.1.** (i)  $(\forall n \geq 0)$  the canonical map

$$S(E/K_n) \longrightarrow (S_\infty)^{\Gamma_n}$$

has finite kernel and cokernel.

(ii) There is an isomorphism of  $\Lambda$ -modules (of finite type)

$$X_\infty \xrightarrow{\sim} \text{Hom}_\Lambda(\widehat{S_\infty}, \Lambda)$$

(where  $\widehat{M}$  denotes the Pontryagin dual of  $M$ ), hence  $\text{rk}_\Lambda(X_\infty) = \text{corank}_\Lambda(S_\infty)$ .

(iii)  $(\forall n \geq 0)$  the canonical map

$$(X_\infty)_{\Gamma_n} \longrightarrow X(E/K_n)$$

has finite kernel.

(iv)  $X_\infty \xrightarrow{\sim} \Lambda^r$  for some  $r \geq 0$ .

(v)  $E(K_\infty)_{\text{tors}}$  is finite.

*Proof.* (i) [Man], Thm. 4.5 or [Maz1], Prop. 6.4 (note that [Man], Lemma 4.6 eliminates the need for the second assumption in [Maz1], 6.1; however, the latter is satisfied in our situation, because of (Heeg)). (ii) [PR], Lemma 5, p. 417. (iii) This follows from (i) and (ii) (cf. [PR], Lemma 4, p. 415). (iv) The R.H.S. in (ii) is reflexive, hence free over  $\Lambda$ . (v) [NeSc], 2.2.

**Lemma 2.2.**  $y \neq 0$  in  $X_\infty$ .

*Proof.* If  $y_n = 0$  in  $E(K_n) \otimes \mathbf{Z}_p$  for all  $n > n_0$ , then

$$x_n \otimes \alpha = x_{n-1} \otimes 1$$

in  $E(K_n) \otimes \mathbf{Q}_p$  for all  $n > n_0$ . As both  $x_n \otimes 1$  and  $x_{n-1} \otimes 1$  lie in  $E(K_n) \otimes \mathbf{Q}$ , but  $\alpha \notin \mathbf{Q}$ , we must have  $x_n \otimes 1 = 0$  in  $E(K_n) \otimes \mathbf{Q}$  for all  $n > n_0$ , which contradicts (now proven) Mazur's conjecture 1.4.

**Lemma 2.3.**  $X_\infty \xrightarrow{\sim} \Lambda$ .

*Proof.* (Note that the implication [Mazur's conjecture  $\implies X_\infty \xrightarrow{\sim} \Lambda$ ] was proved under more restrictive assumptions by Bertolini [Be]).

By Lemma 2.2 there is an exact sequence of  $\Lambda$ -modules of finite type

$$0 \longrightarrow \Lambda y \longrightarrow X_\infty \longrightarrow X_\infty / \Lambda y \longrightarrow 0.$$

For every surjective character

$$\chi : \Gamma \longrightarrow \mu_{p^n}$$

we put  $c(\chi) = n$  and denote by

$$e_\chi = \frac{1}{p^n} \sum_{g \in \Gamma / \Gamma_n} \chi(g)^{-1} g \in \mathbf{Q}_p(\mu_{p^n})[\Gamma / \Gamma_n]$$

the corresponding idempotent. Denote by  $A$  the (finite) set of integers  $n \geq 1$  such that

$$\frac{\omega_n}{\omega_{n-1}} \in \text{Supp}_\Lambda((X_\infty / \Lambda y)_{\text{tors}})$$

(where  $\omega_n = \gamma^{p^n} - 1$ , as usual). If  $c(\chi) \notin A$ , then

$$e_\chi(\mathbf{Q}_p(\mu_{p^n}) \cdot y \bmod \omega_n X_\infty) \neq 0$$

in  $(X_\infty)_{\Gamma_n} \otimes_{\mathbf{Z}_p} \mathbf{Q}_p(\mu_{p^n})$ , hence also

$$e_\chi(\mathbf{Q}_p(\mu_{p^n}) \cdot y_n) \neq 0$$

in  $X(E/K_n) \otimes_{\mathbf{Z}_p} \mathbf{Q}_p(\mu_{p^n})$ , by Lemma 2.1(iii).

According to a mild generalization of the main result of [Be-Da],

$$e_\chi(\mathbf{Q}_p(\mu_{p^n}) \cdot y_n) \neq 0 \implies e_\chi(\mathbf{Q}_p(\mu_{p^n}) \cdot y_n) = e_\chi(X(E/K_n) \otimes_{\mathbf{Z}_p} \mathbf{Q}_p(\mu_{p^n})).$$

Putting everything together and again appealing to Lemma 2.1(iii), we see that

$$\text{rk}_{\mathbf{Z}_p}((X_\infty)_{\Gamma_n}) \leq p^n + O(1),$$

hence  $r \leq 1$  in Lemma 2.1(iv). However,  $r \neq 0$  by Lemma 2.2.

**2.4** Recall that finite  $\Lambda$ -modules are also called *pseudo-null*. Denote by  $(\Lambda\text{Mod})$  the category of all  $\Lambda$ -modules and by  $(\Lambda\text{Mod})/(ps - null)$  the category obtained from  $(\Lambda\text{Mod})$  by inverting all morphisms which have pseudo-null both kernel and cokernel. This is again an abelian category.

**Lemma 2.5.** *In  $(\Lambda\text{Mod})/(ps - \text{null})$  there is an exact sequence*

$$0 \longrightarrow Y \oplus Y \oplus Z \longrightarrow \widehat{S_\infty} \longrightarrow \Lambda \longrightarrow 0$$

*and an isomorphism*

$$Z \xrightarrow{\sim} \bigoplus_{i=1}^k (\Lambda/p^{m_i}\Lambda).$$

*Proof.* Combining Lemma 2.3 with Lemma 2.1(ii) we get an exact sequence in  $(\Lambda\text{Mod})/(ps - \text{null})$

$$0 \longrightarrow (\widehat{S_\infty})_{\text{tors}} \longrightarrow \widehat{S_\infty} \longrightarrow \Lambda \longrightarrow 0.$$

The duality results of [Ne] imply that

$$(\widehat{S_\infty})_{\text{tors}} \xrightarrow{\sim} Y \oplus Y \oplus Z$$

in  $(\Lambda\text{Mod})/(ps - \text{null})$ , with  $Z$  as in the statement of the Lemma (in fact, the condition (Heeg) implies that  $Z$  itself is pseudo-null, at least if  $p > 2$ ; however, we do not need this fact).

### 3. Main Results

**Theorem B.** *Under the assumptions (Heeg) and  $p \nmid Na_p$  we have*

$$s_p(E/K) \equiv 1 \equiv r_{an}(E/K) \pmod{2}.$$

*Proof.* First of all, (Heeg) implies that  $r_{an}(E/K)$  is odd. Lemma 2.5 together with Lemma 2.1(i) give an exact sequence

$$0 \longrightarrow (Y_\Gamma \oplus Y_\Gamma) \otimes_{\mathbf{Z}_p} \mathbf{Q}_p \longrightarrow \text{Hom}_{\mathbf{Z}_p}(X(E/K), \mathbf{Q}_p) \longrightarrow \mathbf{Q}_p \longrightarrow 0,$$

hence

$$s_p(E/K) = 1 + 2 \text{rk}_{\mathbf{Z}_p}(Y_\Gamma) \equiv 1 \pmod{2}.$$

**Theorem A.** *Let  $E$  be an elliptic curve over  $\mathbf{Q}$  with good ordinary reduction at  $p$ . Then*

$$r_{an}(E/\mathbf{Q}) \equiv s_p(E/\mathbf{Q}) \pmod{2}.$$

*Proof.* For every  $K = \mathbf{Q}(\sqrt{D})$  satisfying (Heeg) we have

$$\begin{aligned} s_p(E/K) &= s_p(E/\mathbf{Q}) + s_p(E_D/\mathbf{Q}) \\ r_{an}(E/K) &= r_{an}(E/\mathbf{Q}) + r_{an}(E_D/\mathbf{Q}), \end{aligned}$$

where

$$E_D : Dy^2 = x^3 + Ax + B$$

is the quadratic twist of  $E$  over  $K$ . We distinguish two cases:

(I)  $r_{an}(E/\mathbf{Q})$  is odd.

According to [Wa] there exists  $K$  satisfying (Heeg) such that

$$r_{an}(E_D/\mathbf{Q}) = 0.$$

Results of Kolyvagin [Ko] then imply  $s_p(E_D/\mathbf{Q}) = 0$ , hence

$$s_p(E/\mathbf{Q}) = s_p(E/K) \equiv r_{an}(E/K) = r_{an}(E/\mathbf{Q}) \pmod{2}$$

by Theorem B.

(II)  $r_{an}(E/\mathbf{Q})$  is even.

Choose any  $K$  satisfying (Heeg) and  $p \nmid D$ . Applying the result of Case (I) to  $E_D$  (which has good ordinary reduction at  $p$ ), we obtain

$$s_p(E_D/\mathbf{Q}) \equiv r_{an}(E_D/\mathbf{Q}) \pmod{2},$$

hence

$$s_p(E/\mathbf{Q}) \equiv 1 - s_p(E_D/\mathbf{Q}) \equiv 1 - r_{an}(E_D/\mathbf{Q}) \equiv r_{an}(E/\mathbf{Q}) \pmod{2},$$

again using Theorem B.

#### 4. Higher dimensional quotients of $J_0(N)$

Results of Section 3 can be generalized as follows.

**4.1** Let  $f = \sum_{n \geq 1} a_n q^n \in S_2(\Gamma_0(N))$  be a normalized ( $a_1 = 1$ ) newform. Put  $F(f) = \mathbf{Q}(a_1, a_2, \dots)$ ; this is a totally real number field. Let  $A$  be a quotient abelian variety of  $J_0(N)$  corresponding to  $f$ ; it has dimension  $[F(f) : \mathbf{Q}]$ , is defined over  $\mathbf{Q}$  and is unique up to isogeny. One has an embedding  $\iota : F(f) \hookrightarrow \text{End}(A) \otimes \mathbf{Q}$ ; for each  $n \geq 1$ ,  $\iota(a_n)$  is induced by the Hecke correspondence  $T(n) : J_0(N) \rightarrow J_0(N)$  (with respect to the Albanese functoriality).

**4.2** For each prime number  $p$ ,  $V_p(A) = T_p(A) \otimes_{\mathbf{Z}_p} \mathbf{Q}_p$  is a free module of rank 2 over  $F(f) \otimes_{\mathbf{Q}} \mathbf{Q}_p = \bigoplus_{\mathfrak{p}|p} F(f)_{\mathfrak{p}}$ . Fix a prime  $\mathfrak{p}$  above  $p$  in  $F(f)$ ; then the  $F(f)_{\mathfrak{p}}$ -component  $V_{\mathfrak{p}}(A)$  of  $V_p(A)$  defines a two-dimensional Galois representation

$$G_{\mathbf{Q}} = G(\overline{\mathbf{Q}}/\mathbf{Q}) \longrightarrow \text{Aut}_{F(f)_{\mathfrak{p}}}(V_{\mathfrak{p}}(A)) \xrightarrow{\sim} GL_2(F(f)_{\mathfrak{p}})$$

which is unramified outside  $Np$  and satisfies

$$\det(1 - X \cdot Fr_{\text{geom}}(\ell)|V_{\mathfrak{p}}(A)) = 1 - a_{\ell}\ell^{-1}X + \ell^{-1}X^2$$

for all prime numbers  $\ell \nmid Np$ .

**4.3** Let  $\mathcal{O}_{\mathfrak{p}}$  be the ring of integers of  $F(f)_{\mathfrak{p}}$ . For every number field  $F$ , denote by  $S_{\mathfrak{p}}(A/F)$  the  $\mathfrak{p}$ -primary part of the Selmer group  $S_p(A/F) = \varprojlim_n S(A/F, p^n)$ . This is an  $\mathcal{O}_{\mathfrak{p}}$ -module of cofinite type; denote by  $s_{\mathfrak{p}}(A/F)$

its  $\mathcal{O}_{\mathfrak{p}}$ -corank. Note that  $S_{\mathfrak{p}}(A/F)$  depends only on the  $G(\overline{\mathbf{Q}}/F)$ -module  $V_{\mathfrak{p}}(A)/T_{\mathfrak{p}}(A)$ , where  $T_{\mathfrak{p}}(A)$  is the  $\mathcal{O}_{\mathfrak{p}}$ -component of  $T_p(A)$ , as it coincides with a Bloch-Kato Selmer group

$$S_{\mathfrak{p}}(A/F) = H_f^1(F, V_{\mathfrak{p}}(A)/T_{\mathfrak{p}}(A)).$$

The role of  $r_{an}(E/F)$  is played by the order of vanishing

$$r_{an}(f, F) = \text{ord}_{s=1} L(f \otimes F, s).$$

Under the assumptions (Heeg), (G) and

$$(Ord') \quad \text{ord}_{\mathfrak{p}}(a_p) = 0,$$

the arguments in Section 2 go through for the  $\mathcal{O}_{\mathfrak{p}}$ -modules  $S_{\mathfrak{p}}(A/K_n)$  (Mazur's control Theorem has to be replaced by a purely cohomological "control theorem"; see [Gre]). Similarly, all of the arguments of Section 3 work, if we replace the reference [Ko] by [KoLo]. The final results are the following.

**Theorem B'.** Under the assumptions (Heeg),  $p \nmid N$  and  $\text{ord}_{\mathfrak{p}}(a_p) = 0$  we have

$$s_{\mathfrak{p}}(A/K) \equiv 1 \equiv r_{an}(f, K) \pmod{2}.$$

**Theorem A'.** Assume that  $p \nmid N$  and  $\text{ord}_{\mathfrak{p}}(a_p) = 0$ . Then

$$\text{ord}_{s=1} L(f, s) \equiv s_{\mathfrak{p}}(A/\mathbf{Q}) \pmod{2}.$$

More general results can be deduced by applying the techniques of [NePl]. This will be discussed in a separate publication.

## References

- [Be] M. Bertolini, *Selmer groups and Heegner points in anticyclotomic  $\mathbf{Z}_p$ -extensions*, Compositio Math. **99** (1995), no. 2, 153–182.
- [BeDa] M. Bertolini, H. Darmon, *Kolyvagin's descent and Mordell-Weil groups over ring class fields*, J. Reine Angew. Math. **412** (1990), 63–74.
- [Bi] B. Birch, *Heegner points of elliptic curves*, In: Symposia Mathematica, Vol. XV (Convegno di Strutture in Corpi Algebrici, INDAM, Rome, 1973), pp. 441–445. Academic Press, London, 1975.
- [BCDT] C. Breuil, B. Conrad, F. Diamond, R. Taylor, *On the modularity of elliptic curves over  $\mathbf{Q}$* , preprint.
- [Co] C. Cornut, *Mazur's conjecture on higher Heegner points*, preprint.
- [Gre] R. Greenberg, *Iwasawa theory for  $p$ -adic representations*, In: Algebraic number theory, 97–137, Adv. Stud. Pure Math. **17**, Academic Press, Boston, MA, 1989.
- [Gro] B.H. Gross, *Heegner points on  $X_0(N)$* , In: Modular forms (Durham, 1983), 87–105, Ellis Horwood Ser. Math. Appl., Horwood, Chichester, 1984.
- [Ko] V.A. Kolyvagin, *Euler systems*, In: The Grothendieck Festschrift, Vol. II, 435–483, Progress in Math. **87**, Birkhäuser Boston, Boston, MA, 1990.
- [KoLo] V.A. Kolyvagin, D.Yu. Logachev, *Finiteness of the Shafarevich-Tate group and the group of rational points for some modular abelian varieties* (Russian), Algebra i Analiz **1** (1989), No. 5, 171–196. English translation: Leningrad Math. J. **1** (1990), No. 5, 1229–1253.
- [Man] Yu.I. Manin, *Cyclotomic fields and modular curves*, Uspehi Mat. Nauk **26** (1971), no. 6(162), 7–71. (Russian)
- [Maz1] B. Mazur, *Rational points of abelian varieties with values in towers of number fields*, Invent. Math. **18** (1972), 183–266.
- [Maz2] B. Mazur, *Modular curves and arithmetic*, In: Proceedings of the ICM 1983 (Warsaw), Vol. 1, 185–211, PWN, Warsaw, 1984.
- [Ne] J. Nekovář, *Selmer complexes*, in preparation.
- [NePl] J. Nekovář, A. Plater, *On the parity of ranks of Selmer groups*, Asian J. Math. **4** (2000), No. 2, 437–498.
- [NeSc] J. Nekovář, N. Schappacher, *On the asymptotic behaviour of Heegner points*, Turkish J. of Math. **23** (1999), No. 4, 549–556.
- [PR] B. Perrin-Riou, *Fonctions  $L$   $p$ -adiques, théorie d'Iwasawa et points de Heegner*, Bull. Soc. Math. France **115** (1987), No. 4, 399–456.
- [Va1] V. Vatsal, *Uniform distribution of Heegner points*, preprint.
- [Va2] V. Vatsal, *Special values of anticyclotomic  $L$ -functions*, preprint.
- [Wa] J.-L. Waldspurger, *Correspondances de Shimura*, In: Proceedings of the ICM 1983 (Warsaw), Vol. 1, 525–531, PWN, Warsaw, 1984.

Department of Pure Mathematics and Mathematical Statistics  
University of Cambridge  
Centre for Mathematical Sciences  
Wilberforce Road  
Cambridge CB3 0WB  
UK  
nekovar@dpmms.cam.ac.uk